

William's E, powder, without L-glutamine

Cat. No.: PM151211P

Size: 5×1L / 1×10L / 1×50L / 100L / 500L

General Information

Product Form	Powder
D-Glucose	2000 mg/L
Concentration	10.48 g/L
HEPES	Negative
L-Glutamine	Negative
NaHCO ₃	Negative
Phenol red	10 mg/L
Sodium pyruvate	0.23 mM
Storage	2-8°C, Shading Light
Shipping	Room Temperature
Expiration date	36 months

Background

William's E medium was designed by Willam and Gunn, which is mainly used for long-term culture of rat liver epithelial cells, and can also be used for the culture of hepatocytes from other prolactinuous animals. This product contains many kinds of cell culture required amino acids, vitamins, inorganic salts and other ingredients, but does not contain protein, lipids and any growth factors, so this product needs to be used with serum or serum-free additives.

Preparation method

1. The preparation water should be purified water, ultra-pure water or water for injection (WFI), and the water temperature should be controlled between 20-30°C during the preparation process.
2. Measure 90% of the final volume preparation water to the solution preparation system. Start stirring, and avoid generating bubbles. For example, if 1L is required, add 900 mL of preparation water here. And it's recommended that the power output per unit volume (P/V) of the mixing system is greater than 10 W/m³.
3. Weigh the appropriate amount of powder according to the concentration of 10.48 g/L accurately, and add it to the container prepared in step 2. Stir for more than 20 minutes dissolve all powder completely.
4. After the solution is clear, add NaHCO₃ at a concentration of 2.2 g/L, continue stirring for 5-10 minutes until dissolved, then add ultra pure water to adjust the volume to the 100% of required.
5. If necessary, adjust the pH to 7.20-7.30 with 1 mol/L NaOH solution or 1 mol/L HCl solution. Since filtration will slightly increase the pH, the pH value here is lower than the target pH value (7.20-7.40).
6. The prepared solution should be sterilized using a 0.2 µm pore size filter membrane under positive pressure (ensure aseptic technique).

7. After filtration, a small amount of liquid culture medium can be taken for quality inspection, and use only after passing the test.
8. The filtered liquid medium should be used immediately or stored in glass bottles, culture medium bottles (PET), or single-use storage bags with an oxygen-barrier coating at 2-8°C away from light. The liquid medium has a shelf life of 2 year under these conditions.

Notes

1. This product is only used for scientific research or further research, not for diagnosis and treatment.
2. Please wear a lab coat and use disposable gloves and a mask during operation.
3. To ensure the optimal performance of this product, please strictly adhere to the recommended storage conditions for its preservation.