BVES / POPDC1 Antibody, Rabbit PAb, Antigen Affinity Purified


Catalog Number: **101360-T40**

GENERAL INFORMATION	
Immunogen:	A synthetic peptide corresponding to the C-terminus of the Human BVES / POPDC1
Preparation	Produced in rabbits immunized with a synthetic peptide corresponding to the C-terminus of the Human BVES / POPDC1, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human BVES / POPDC1
Formulation:	0.2 μm filtered solution in PBS
Storage:	This antibody can be stored at 2℃-8℃ for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20℃ to -80℃. Preservative-Free. Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.
APPLICATIONS	
Applications:	WB,IHC-P
RECOMMENDED CONCENTRATION	
IHC-P	IHC-P: 1:500-1:2000
Western Blot	WB: 1:500-1:2000

Please Note: Optimal concentrations/dilutions should be determined by the end user.


BVES / POPDC1 Antibody, Rabbit PAb, Antigen Affinity Purified

Catalog Number: **101360-T40**



Immunochemical staining of human BVES in human skeletal muscle with rabbit polyclonal antibody (1:1000, formalin-fixed paraffin embedded sections).

(Validation Experiment)




Anti-BVES rabbit polyclonal antibody at 1:500 dilution.
Lane A: Mosue skeletal muscle tissue lysate
Lane B: 293 Whole Cell Lysate

Lysates/proteins at 30 µg per lane.
Secondary
Goat Anti- Rabbit IgG H&L (Dylight 800) at 1/10000 dilution.

Developed using the Odyssey technique.
Performed under reducing conditions.

Predicted band size:41 kDa
Observed band size:40 kDa
(We are unsure as to the identity of these extra bands.)

(Validation Experiment)



Anti-BVES rabbit polyclonal antibody at 1:500 dilution
Lane A: BVES konckout Hela Whole Cell Lysate
Lane B: Hela Whole Cell Lysate

Lysates/proteins at 10 µg per lane.
Secondary
Goat Anti- Rabbit IgG H&L (Dylight 800) at 1/10000 dilution.

Developed using the ECL technique.
Performed under reducing conditions.

Predicted band size:41 kDa
Observed band size:40 kDa
(We are unsure as to the identity of these extra bands.)

(Validation Experiment)