IL-1F6 Antibody, Rabbit PAb, Antigen Affinity Purified


Catalog Number: **10607-T52**

GENERAL INFORMATION	
Immunogen:	Recombinant Human IL-1F6 Protein (Catalog#10607-HNAE)
Preparation	Produced in rabbits immunized with purified, recombinant Human IL-1F6 (rh IL-1F6; Catalog#10607-HNAE; Q9UHA7; Met1-Phe158). IL-1F6 specific IgG was purified by Human IL-1F6 affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human IL-1F6
Formulation:	0.2 μm filtered solution in PBS
Storage:	This antibody can be stored at 2℃-8℃ for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20℃ to -80℃. Preservative-Free. Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.
APPLICATIONS	
Applications:	WB,ELISA,IP
RECOMMENDED CONCENTRATION	
Western Blot	WB: 1:500-1:2000
Immunoprecipitation	IP: 5-10 μL/mg of lysate
ELISA	ELISA: 1:5000-1:10000 This antibody can be used at 1:5000-1:10000 with the appropriate secondary reagents to detect Human IL-1F6.

Please Note: Optimal concentrations/dilutions should be determined by the end user.

IL-1F6 Antibody, Rabbit PAb, Antigen Affinity Purified

Catalog Number: **10607-T52**




Anti-IL1F6 rabbit polyclonal antibody at 1:500 dilution
Lane A: MDA-MB231 Whole Cell Lysate
Lane B: HT1080 Whole Cell Lysate
Lane C: MCF7 Whole Cell Lysate

Lysates/proteins at 30 µg per lane.
Secondary
Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique.
Performed under reducing conditions.

Predicted band size:18 kDa
Observed band size:24 kDa



IL1F6 was immunoprecipitated using:
Lane A:0.5 mg MDA-MB231 Whole Cell Lysate

2 µL anti-IL1F6 rabbit polyclonal antibody and 60 µg of Immunomagnetic beads Protein A/G.

Primary antibody:
Anti-IL1F6 rabbit polyclonal antibody,at 1:100 dilution

Secondary antibody:
Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution

Developed using the ECL technique.
Performed under reducing conditions.

Predicted band size: 18 kDa
Observed band size :23 kDa