MID1IP1 Antibody, Rabbit PAb, Antigen Affinity Purified


Catalog Number: 201198-T36

GENERAL INFORMATION	
Immunogen:	E. coli-derived Human MID1IP1 fragment
Preparation	Produced in rabbits immunized with E. coli-derived Human MID1IP1 fragment, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human MID1IP1
Formulation:	PBS, pH7.0 with 0.03% Proclin300
Storage:	This antibody can be stored at 2℃-8℃ for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20℃ to -80℃. Avoid repeated freeze-thaw cycles.
Alternative Names:	MID1IP1
APPLICATIONS	
Applications:	WB, IP
RECOMMENDED CONCENTRATION	
Western Blot	WB: 1:500-1:2000
Immunoprecipitation	IP:1-5μL/mg of lysate

Please Note: Optimal concentrations/dilutions should be determined by the end user.

MID1IP1 Antibody, Rabbit PAb, Antigen Affinity Purified

Catalog Number: **201198-T36**




Anti-MID1IP1 rabbit polyclonal antibody at 1:500 dilution
Lane A: HepG2 Whole Cell Lysate
Lane B: HEK293 Whole Cell Lysate

Lysates/proteins at 30 µg per lane.
Secondary
Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique.
Performed under reducing conditions.

Predicted band size:20 kDa
Observed band size:25 kDa
(We are unsure as to the identity of these extra bands.)



MID1IP1 was immunoprecipitated using:
Lane A:0.5 mg HepG2 Whole Cell Lysate

4 µL anti-MID1IP1 rabbit polyclonal antibody and 60 µg of Immunomagnetic beads Protein A/G.

Primary antibody:
Anti-MID1IP1 rabbit polyclonal antibody,at 1:100 dilution

Secondary antibody:
Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution

Developed using the ECL technique.
Performed under reducing conditions.

Predicted band size: 20 kDa
Observed band size :25 kDa