CD2BP2 Antibody, Rabbit PAb, Antigen Affinity Purified


Catalog Number: **201328-T36**

GENERAL INFORMATION	
Immunogen:	E. coli-derived Human CD2BP2 fragment
Preparation	Produced in rabbits immunized with E. coli-derived Human CD2BP2 fragment, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human CD2BP2
Formulation:	PBS, pH7.0 with 0.03% Proclin300
Storage:	This antibody can be stored at 2℃-8℃ for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20℃ to -80℃. Avoid repeated freeze-thaw cycles.
Alternative Names:	CD2BP2
APPLICATIONS	
Applications:	WB, IP
RECOMMENDED CONCENTRATION	
Western Blot	WB: 1:500-1:2000
Immunoprecipitation	IP:1-5μL/mg of lysate

Please Note: Optimal concentrations/dilutions should be determined by the end user.

CD2BP2 Antibody, Rabbit PAb, Antigen Affinity Purified

Catalog Number: **201328-T36**




Anti-CD2BP2 rabbit polyclonal antibody at 1:500 dilution
Lane A: HeLa Whole Cell Lysate
Lane B: Jurkat Whole Cell Lysate
Lane C: U-251MG Whole Cell Lysate

Lysates/proteins at 30 µg per lane.
Secondary
Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique.
Performed under reducing conditions.

Predicted band size:38 kDa
Observed band size:55 kDa
(We are unsure as to the identity of these extra bands.)



CD2BP2 was immunoprecipitated using:
Lane A:0.5 mg Jurkat Whole Cell Lysate

4 µL anti-CD2BP2 rabbit polyclonal antibody and 60 µg of Immunomagnetic beads Protein A/G.

Primary antibody:
Anti-CD2BP2 rabbit polyclonal antibody,at 1:100 dilution

Secondary antibody:
Clean-Blot IP Detection Reagent (HRP) at 1:1000dilution

Developed using the ECL technique.
Performed under reducing conditions.

Predicted band size: 51 kDa
Observed band size :55 kDa