

DDX21 Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological
Biological Solution Specialist

Catalog Number: 202536-T38

GENERAL INFORMATION	
Immunogen:	E. coli-derived Human DDX21 fragment
Preparation	Produced in rabbits immunized with E. coli-derived Human DDX21 fragment, and purified by antigen affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human DDX21
Formulation:	PBS, pH7.0 with 0.03% Proclin300
Storage:	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Alternative Names:	DDX21, GUA, GURDB, RH-II/GU, RH-II/GuA
APPLICATIONS	
Applications:	WB, ICC/IF, IP
RECOMMENDED CONCENTRATION	
ICC/IF	ICC/IF: 1:100-1:500
Western Blot	WB: 1:500-1:2000
Immunoprecipitation	IP: 0.5-2µL/mg of lysate


Please Note: Optimal concentrations/dilutions should be determined by the end user.

DDX21 Antibody, Rabbit PAb, Antigen Affinity Purified


Catalog Number: 202536-T38



Sino Biological
Biological Solution Specialist



Immunofluorescence staining of DDX21 in U251MG cells. Cells were fixed with 4% PFA, permeabilized with 0.1% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human DDX21 polyclonal antibody (dilution ratio 1:200) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to Nucleus.



Anti-DDX21 rabbit polyclonal antibody at 1:500 dilution

Lane A: HepG2 Whole Cell Lysate
Lane B: HEK293 Whole Cell Lysate
Lane C: Jurkat Whole Cell Lysate
Lane D: HeLa Whole Cell Lysate

Lysates/proteins at 30 µg per lane.

Secondary
Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique. Performed under reducing conditions.

Predicted band size: 87 kDa
Observed band size: 95 kDa



IARS2 was immunoprecipitated using:
Lane A: 0.5 mg HepG2 Whole Cell Lysate

4 µL anti-IARS2 rabbit polyclonal antibody and 60 µg of Immunomagnetic beads Protein A/G. Primary antibody:

Anti-IARS2 rabbit polyclonal antibody, at 1:100 dilution

Secondary antibody:
Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution

Developed using the ECL technique. Performed under reducing conditions.

Predicted band size: 87 kDa
Observed band size: 95 kDa